

Role of Zinc Deficiency in Photosynthetic Pigments and Peroxidase Activity of Tomato Seedlings

Taylan Kösesakal* and Muammer Ünal

Istanbul University, Faculty of Science, Department of Biology,
34134 Suleymaniye, Istanbul / Turkey

Abstract

Zinc (Zn) deficiency is occurring in different climate regions and almost in all countries. Zinc deficiency is a common micronutrient deficiency in plants and causes severe reductions in crop production. In this study the effect of zinc deficiency on pigment content and peroxidase activity in early growth phase of tomato seedlings was investigated. For this purpose, Hoagland solutions with or without zinc ($ZnCl_2$) were used as growth medium to determine the possible effects of zinc deficiency. In order to elucidate the physiological consequences of zinc deficiency; anthocyanin, carotenoid, and chlorophyll content and peroxidase enzyme activity were measured in roots, hypocotyls, and cotyledons of 8 day-old plants. Zn deficiency promoted anthocyanin and carotenoid accumulation in hypocotyl and cotyledon tissues of tomato seedlings. Furthermore, Zn deficiency enhanced peroxidase activity especially in the root and hypocotyl tissues of tomato seedlings.

Keywords: Chlorophyll, Peroxidase, Tomato, Zinc, Zinc deficiency

***Corresponding author:** Taylan Kösesakal (E-mail: taylank@istanbul.edu.tr)

(Received: 03.03.2008 Accepted: 18.03.2009)

Introduction

Zinc (Zn) is one of the essential micronutrients playing a significant role in many vital metabolic processes (Rout and Das 2003; Aravind and Prasad 2005a, b). For instance, Zn is a cofactor for several enzymes such as anhydrases (Aravind and Prasad 2004a; 2005c), dehydrogenases, oxidases and peroxidases (Vallee and Auld 1990; Aravind and Prasad 2003). Furthermore, Zn could be also involved in chlorophyll formation by taking part in the regulation of cytoplasmic concentrations of nutrients. For example, Fischer (1997) reported that Mg-deficiency decreased chlorophyll concentration in beans, and Welkie et al. (1990) recorded that there was a linear relationship between leaf-Fe and chlorophyll content in peppers. Zn could also increase the biosynthesis of chlorophyll and carotenoids ultimately proving beneficial for the photosynthetic machinery of the plant system (Aravind and Prasad 2004b).

Conclusively, Zn is a must for plant development. Zinc deficiency is a major global problem hindering plant cultivation, and this problem is especially exacerbated in acidic calcareous soils which is the most common soil type in arid and semi-arid regions of the world (Cakmak 2000; Hacısalihoglu et al. 2004). Zinc deficiency reduces plant growth and inhibits photosynthesis in a wide variety of plants including maize (Wang and Jin 2005) and rice (Wenrong et al. 2008). Since the most parts of Turkey lie in arid or semi-arid climatic zones, zinc deficiency is one of the most serious problems restricting plant yield, particularly in Central Anatolia. Although Turkish soils are rich in total Zn content, the bioavailability of Zn is extremely low due to the highly acidic nature of soils and results in stunted plant growth and reduced yields (Eyüpoğlu et al. 1994; Gülser et al. 2004).

Peroxidases have various physiological roles in plant cells and participate in many reactions

including lignification, cross linking of cell elongation and phenol oxidation, all linked to growth reductions (Mocquot et al. 1996). Peroxidase activity shows a close correlation with changes in physiological processes such as, respiration, photosynthesis and transpiration, and therefore has the potential to serve as a sensitive indicator of compromised metabolic activity (Verkleij and Schat 1990; Aravind and Prasad 2005a). Inhibition of photosynthesis by some stressors in higher plants is well documented. Reduction in the levels of photosynthetic pigments, including chlorophylls a and b and accessory pigments such as carotenoids, on exposure to biotic or abiotic stressors have been observed in many species (Macfarlane and Burchett 2001; Thao and Yanyun 2005; Lau et al. 2006). The inhibition of both photosystem II and photosystem I generating lipid peroxidation from active oxygen radicals, may also result in the destruction of photosynthetic pigments and thylakoid membrane structure (Droppa and Horvath 1990). Anthocyanins are best known as the characteristic red, blue, and purple of plant tissues. They belong to the widespread class of phenolic compounds collectively named flavonoids, produced in the cytoplasm and then transported into the vacuole (Harborne 1988; Shirley 1996). The synthesis of anthocyanins is induced by UV-B (Wilson et al. 1998, Warren et al. 2003), nutrient deficiency (Pinto et al. 1999), low temperature (Rabino and Mancinelli 1986), water deficit (Nogués et al. 1998), and heavy metal stress (Ling-Peng et al. 2006). Thus, both peroxidases and pigments may have the potential to be employed as sensitive indicators of biotic or abiotic stressors and may predict subsequent events at the organism level (Dietz et al. 1999).

The main objective of this study was to investigate the effects of zinc deficiency on pigment content and peroxidase activity in the early growth phase of tomato seedlings. In this study, we determined quantitatively the change in chlorophyll, carotenoid, anthocyanin, and

peroxidase activity which will be expected to shed light on the role of pigment in tomato seedlings and zinc deficiency.

Material and Methods

Tomato plants (*Lycopersicon esculentum* Mill.) were grown in standard Hoagland and Arnon (1938) medium, with Zn [+Zn, ZnCl₂, control] and without Zn [-Zn, treatment]. After sterilization with 70 % ethanol, tomato seeds were imbibed at room temperature for 24 hours. Subsequently, the seeds were washed with distilled water and transferred into petri dishes and wrapped into the filter papers wetted with appropriate nutrient solutions (+Zn and -Zn). Next, each petri dish was watered with distilled water for every two days. Experimental plants were grown for 12 hours of light and 12 hours of dark cycles in 25 ±1 °C and 8000 lux light intensity. For analyses, roots, hypocotyls and cotyledons of the plants were collected at appropriate times.

After the 8 days of growth under the above mentioned conditions the same method was employed as in Manchinelli et al. (1990) for the determination of anthocyanin amounts in the cotyledons and hypocotyls of 8 day-old seedlings. For the determination of chlorophyll and carotenoid amounts in the same tissues, the same approach was utilized as explained by Parsons and Strickland (1963). Finally, peroxidase enzyme activity in the roots, cotyledons and hypocotyls were calculated by the absorbance ratios at 470 nm in 0.1 mM phosphate buffer (pH 5.8), 15 mM guaiacol and 5 mM H₂O₂ solutions (Birecka et al. 1973).

One-way ANOVA test was employed for statistical analyses of the obtained data (P<0.01). Vertical bars show standard errors.

Results

The anthocyanin content in the hypocotyls and cotyledons of the tomato seedlings grown in the medium without ZnCl₂ was much higher than the control (Fig. 1).




Figure 1. The amount of anthocyanin, hypocotyls, and cotyledons of 8 day old tomato seedlings grown in normal Hoagland solution containing zinc (control) and without zinc (-Zn).

As for chlorophyll, it was observed that the amount of chlorophyll in the hypocotyls of seedlings grown in -Zn increased compared to the control plants. When we compared the level of chlorophyll in the cotyledon tissues of plants

grown in control medium and in Hoagland solution without -Zn, the total amount of chlorophyll was almost equal, but there was some difference in the relative ratios of chlorophyll a and chlorophyll b (Table 1 and Fig. 2).

Table 1. Chlorophyll contents ($\mu\text{g/g}$ FW) of hypocotyls, and cotyledons of tomato seedlings grown in normal Hoagland solution containing zinc (control) and without zinc (-Zn).

	Hypocotyl			Cotyledon		
	Chlorophyll a	Chlorophyll b	Total Chlorophyll	Chlorophyll a	Chlorophyll b	Total Chlorophyll
Control	95,2 \pm 6,8	46,7 \pm 5,5	141,9 \pm 9,7	642,1 \pm 11,1	409,4 \pm 16,6	1051,5 \pm 21,9
(-Zn)	112,7 \pm 10,1	57,9 \pm 7,4	170,6 \pm 15,1	575,3 \pm 39,4	477,9 \pm 42,9	1053,2 \pm 30,3




Figure 2. Total chlorophyll contents of hypocotyls, and cotyledons of tomato seedlings grown in normal Hoagland solution containing zinc (control) and without zinc (-Zn).

There was an increase in the carotenoid level of the seedlings grown in Hoagland solutions containing -Zn (Fig. 3).

The peroxidase enzyme activity of tomato seedlings which were grown -Zn was increased

in comparison to the control plants. The peroxidase activity in the root and hypocotyl tissues of tomato seedlings grown in the -Zn medium was noticeably higher than the control (Fig. 4).




Figure 3. Total carotenoid contents of hypocotyls, and cotyledons of tomato seedlings grown in normal Hoagland solution containing zinc (control) and without zinc (-Zn).




Figure 4. Peroxidase activity (ΔA/g FW.min.) of roots, hypocotyls, and cotyledons of tomato seedlings grown in normal Hoagland solution containing zinc (control) and without zinc (-Zn).

Discussion

Anthocyanins have long been considered to act as a screen against UV-B light because the latter induced reddening. More recent studies have found that anthocyanins are produced in response to various types of stress, including metal stress (Chalker-Scott 1999; Hale et al. 2001). In general, anthocyanins are believed to increase the antioxidant response of plants in order to uphold the regular physiological status in tissues directly or indirectly affected by biotic or abiotic stressors (Yamasaki 1996; Rice-Evans et al. 1997; Neill et al. 2002). Gülçin et al. (2005) also reported that total anthocyanins from *Perilla panchensis* had strong antioxidant activity, reducing power, superoxide anion scavenging, hydrogen peroxide scavenging, and metal chelating activities when compared to standard antioxidant compounds such as α -tocopherol and trolox. We observed that the absence of zinc in the medium caused the augmentation of anthocyanin in both hypocotyls and cotyledons of the tomato seedlings. It is a known fact that plants defend themselves against a variety of hostile environmental conditions by inducing the biosynthesis of certain chemicals such as anthocyanins and antioxidants, and by hyping the activity of certain enzymes (Weckx and Clijsters 1997; Mazhoudi et al. 1997). Thus, from the results of our trials, we can claim that zinc deficiency is a stress factor and it increases the anthocyanin contents of tomato seedlings (Fig. 1).

Despite the amount of chlorophyll in the hypocotyl tissues of the tomato seedlings, seedlings grown in the -Zn medium were higher relative to the control, the chlorophyll amount in the cotyledon tissues of these seedlings was not very different than the control but there was some difference in the relative ratios of chlorophyll a and chlorophyll b (Table 1, Fig. 2). Wenrong et al. (2008) reported zinc deficiency caused extensive declines in leaf chlorophyll content and ratios of chl a:b. Also, Hu and Sparks (1991) found that the zinc deficiency reduces the amount of chlorophyll in

the leaves of Stuart pecans. In one other experiment, it is shown that the application of exogenous zinc to leaves of tomato plants caused accumulation of chlorophyll content of leaves at both low and high concentrations (Kaya and Higgs 2002). However, we reached a contradicting result in this experiment; chlorophyll content of cotyledons of the plants grown without zinc was almost same with the values obtained from the control plants (Table 1, Fig. 2).

We denoted that the total carotenoid content of the tomato seedlings grown in media containers without zinc, increased (Fig. 3). It is shown that the decreased carotenoid levels observed in Cd-treated *Ceratophyllum demersum* plants were well maintained with the addition of Zn. (Aravind and Prasad 2004b). Carotenoids are known to quench the oxidizing species and triplet state of the chlorophyll and other excited molecules in the pigment bed, which are seriously involved in disrupting metabolism through oxidative damage to cellular components (Candan and Tarhan 2003). It could be hypothesized that relative change in the carotenoid content in the various tissues could be as result of physiological response caused by zinc deficiency.

It is shown that the peroxidase enzyme activity of tomato seedlings grown without $ZnCl_2$, was increased in comparison to the control plants (Fig. 4). Zinc is known to have a stabilizing and protective effect on the biomembranes against oxidative and peroxidative damage, loss of plasma membrane integrity and also alteration of the permeability of the membrane (Aravind and Prasad 2003). Marschner and Cakmak (1989) had shown an enhancement in the peroxidative damage of crucial cell components (such as membranes, enzymes, etc.) in plants grown under zinc deficiency. Subsequently, Cakmak and Marschner (1993) reported a decline in the activities of various antioxidant enzymes with the exception of guaiacol peroxidase (POD).

The most common plant pigments are chlorophyll, carotenoid, and anthocyanin. The

content of chlorophyll, carotenoid, and anthocyanin as well as their relative amount determine color and appearance of plants (Abbott 1999) and serve as markers of health. In conclusion, our results showed that zinc deficiency induced anthocyanin and carotenoid augmentation, and peroxidase enzyme activity either in hypocotyls or cotyledons of tomato seedlings. Anthocyanin and POD may be involved in internal detoxification mechanisms of the hypocotyls and cotyledons against zinc deficiency.

Acknowledgements

This work was supported by Istanbul University Research Fund (Project No: T-327/03112003). The authors wish to thank Dr Bayram YUKSEL, for his useful comments and grammatical revision of this article.

References

- Abbott J.A. (1999) Quality measurement of fruit and vegetables. *Postharvest Biology and Technology*, 15: 207–225.
- Aravind P. and Prasad M.N.V. (2003) Zinc alleviates cadmium induced toxicity in *Ceratophyllum demersum*, a fresh water macrophyte. *Plant Physiology and Biochemistry*, 41: 391-397.
- Aravind P. and Prasad M.N.V. (2004a) Carbonic anhydrase impairment in cadmium-treated *Ceratophyllum demersum* L.(free floating freshwater macrophyte): toxicity reversal by zinc. *Journal of Analytical Atomic Spectrophotometry*, 19: 52-57.
- Aravind P. and Prasad M.N.V. (2004b). Zinc protects chloroplasts and associated photochemical functions in cadmium exposed *Ceratophyllum demersum* L., a freshwater macrophyte. *Plant Science*, 166: 1321–1327.
- Aravind P. and Prasad M.N.V. (2005a) Modulation of cadmium-induced oxidative stress in *Ceratophyllum demersum* by zinc involves ascorbate-glutathione cycle and glutathione metabolism. *Plant Physiology and Biochemistry*, 43: 107-116.
- Aravind P. and Prasad M.N.V. (2005b) Cadmium-Zinc interactions in hydroponic system using *Ceratophyllum demersum*: adaptive plant ecophysiology, biochemistry and molecular toxicology. *Brazilian Journal of Plant Physiology*, 17 (1): 3-20.
- Aravind P. and Prasad M.N.V. (2005c) Zinc mediated protection to the conformation of carbonic anhydrase in cadmium exposed *Ceratophyllum demersum* L. *Plant Science*, 169: 245-254.
- Birecka H., Briber K.A. and Catalfamo J.L. (1973) Comparative studies on tobacco pit and sweet potato root isoperoxidases in relation to injury, indolacetic acid, and ethylene effects. *Plant Physiology*, 52: 43-49.
- Cakmak I. (2000) Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *New Phytologist*, 146: 185-205.
- Cakmak I. and Marschner H. (1993) Effect of zinc nutritional status on superoxide radical and hydrogen peroxide scavenging enzymes in bean leaves. In: Barrow, N.J. (ed.). *Plant Nutrition—From Genetic Engineering to Field Practice* Kluwer Academic Publishers, Dordrecht, pp. 133-137.
- Candan N. and Tarhan L. (2003) Relationship among chlorophyll-carotenoid content, antioxidant enzyme activities and lipid peroxidation levels by Mg²⁺ deficiency in the *Mentha pulegium* leaves. *Plant Physiology and Biochemistry*, 41: 35-40.
- Chalker-Scott L. (1999) Environmental significance of anthocyanins in plant stress response. *Photochemistry and Photobiology*, 70:1-9.
- Dietz K.J., Baier M. and Kramer U. (1999) Impact of heavy metals on photosynthesis. In M.N.V. Prasad and J. Hagemeyer, *Heavy Metal Stress in Plants: from Molecules to Ecosystems*, eds. pp. 73-97. Springer, Berlin.
- Droppa M. and Horvath G. (1990) The role of Cu in photosynthesis. *Critical Review of Plant Science*, 9 (2): 111-124.

- Eyüpoğlu F., Kurucu N. and Sanısağ U. (1994) Status of plant available micronutrients in Turkish soils. In: Soil and Fertilizer Research Institute Annual Report, No. R-118, Ankara, Turkey (in Turkish), pp. 25-32.
- Fischer, E.S. (1997) Moderate magnesium deficiency affected chlorophyll content of bean plant. *Photosynthetica*, 33 (3-4): 385-390.
- Gülçin I., Berashvili D. and Gepdiremen A. (2005) Antiradical and antioxidant activity of total anthocyanins from *Perilla pankinensis* decne. *Journal of Ethnopharmacology*, 101: 287-293.
- Gülser F., Togay Y. and Togay N. (2004) The Effects of Zinc Application on Zinc Efficiency and Nutrient Composition of Lentil (*Lens culinaris* Medic.) Cultivars. *Pakistan Journal of Biological Sciences*, 7 (5): 751-759.
- Hacısalihoğlu G., Ozturk L., Cakmak I., Welch R. M. and Kochian L. (2004) Genotypic variation in common bean in response to zinc deficiency in calcareous soil. *Plant and Soil*, 259: 71-83.
- Hale K.L., McGrath S.P., Lombi E., Stack S.M., Terry N., Pickering I.J., George G.N. and Pilon-Smits E.A.H. (2001) Molybdenum sequestration in brassica species. A Role for Anthocyanins?. *Plant Physiology*, 126: 1391-1402.
- Harborne J.R. (1988) The flavonoids: Recent advances. In: Goodwin TW, (ed.) *Plant Pigments*. London: Academic Press, pp. 299-343.
- Hoagland D.R. and Arnon D.I. (1938) The water culture method for growing plants without soil. *California Agricultural Experiment Station Circular*, 347: 1-39.
- Hu H. and Sparks D. (1991) Zinc-deficiency inhibits chlorophyll synthesis and gas-exchange in stuart pecan. *Hortscience*, 26: 267-268.
- Kaya C. and Higgs D. (2002) Response of tomato (*Lycopersicon esculentum* L.) cultivars to application of zinc when grown in sand culture at low zinc. *Scientia Horticulturae*, 93: 53-64.
- Lau T.S.L., Eno E., Goldstein G., Smith C. and Christopher D.A. (2006) Ambient levels of UV-B in Hawaii combined with nutrient deficiency decrease photosynthesis in near-isogenic maize lines varying in leaf flavonoids: Flavonoids decrease photoinhibition in plants exposed to UV-B. *Photosynthetica*, 44: 394-403.
- Ling-Peng D., Zhi-Ting X., Yu H. and Min-Jing L. (2006) Cadmium-induced changes in pigments, total phenolics, and phenylalanine ammonia-lyase activity in fronds of *Azolla imbricata*. *Environmental Toxicology*, 21: 505-512.
- Macfarlane G.R. and Burchett M.D., (2001) Photosynthetic pigments and peroxidase activity as indicators of heavy metal stress in the grey mangrove, *Avicennia marina* (Forsk.) Vierh. *Marine Pollution Bulletin*, 42: 233-240.
- Mancinelli A.L. (1990) Interaction between light quality and light quantity in the photoregulation of anthocyanin production. *Plant Physiology*, 92: 1191-1195.
- Marschner H. and Cakmak I. (1989) High light intensity enhances chlorosis and necrosis in leaves of zinc, potassium and magnesium deficient bean (*Phaseolus vulgaris*) plants. *Journal of Plant Physiology*, 134: 308-315.
- Mazhoudi S., Chaoui A., Ghorbal M.H. and El Ferjani E. (1997) Response of antioxidant enzymes to excess copper in tomato (*Lycopersicon esculentum* Mill.). *Plant Science*, 127: 129-137.
- Mocquot B., Vangronsveld J., Clijsters H. and Mench M. (1996) Copper toxicity in young maize (*Zea mays* L.) plants: effects on growth, mineral and chlorophyll contents, and enzyme activities. *Plant and Soil*, 182: 287-300.
- Neill S.O., Gould K.S., Kilmartin P.A., Mitchell K.A. and Markham K.R. (2002) Antioxidant activities of red versus green leaves in *Elatostema rugosum*. *Plant Cell & Environment*, 25: 539-547.

- Nogués S., Allen D.J., Morison J.I.L. and Baker N.R. (1998) Ultraviolet-B radiation effects on water relations, leaf development, and photosynthesis in droughted pea plants. *Plant Physiology*, 117: 173-181.
- Parsons T.R. and Strickland J.D.H. (1963) Discussion of spectrophotometric determination of marine pigments, with revised equations for ascertaining chlorophylls and carotenoids. *Journal of Marine Research*, 21: 115-163.
- Pinto M.E., Casati P., Hsu T.P., Ku M.S.B. and Edwards G.E. (1999) Effects of UV-B radiation on growth, photosynthesis, UV-B absorbing compounds and NADP-malic enzyme in bean (*Phaseolus vulgaris* L.) grown under different nitrogen conditions. *Journal of Photochemistry and Photobiology*, 48: 200-209.
- Rabino I. and Mancinelli A.L. (1986) Light, temperature and anthocyanin production. *Plant Physiology*, 81: 922-924.
- Rice-Evans C.A., Miller N.J. and Paganga G. (1997) Antioxidant properties of phenolic compounds. *Trends in Plant Science*, 2: 152-159.
- Rout G.R. and Das P. (2003) Effect of metal toxicity on plant growth and metabolism. *Agronomie*, 23: 3-11.
- Shirley B.W. (1996) Flavonoid biosynthesis: New functions for an old pathway. *Trends in Plant Science*, 1: 377-382.
- Thao N. and Yanyun Z. (2005). Retaining Green Pigments on Thermally Processed Peels-on Green Pears. *Journal of Food Science*, 70: 568-574.
- Vallee B.L. and Auld D.S. (1990). Zinc coordination, function, and structure of zinc enzymes and other proteins. *Biochemistry*, 29: 5647-5659.
- Verkleij J.A.C. and Schat H. (1990) Mechanism of metal tolerance in plants. In *Heavy Metal Tolerance in Plants-Evolutionary Aspects*, ed. Shaw, A.J., pp.179-193. CRC Press, Boca Raton, FL.
- Wang H. and Jin J.Y. (2005) Photosynthetic rate, chlorophyll fluorescence parameters, and lipid peroxidation of maize leaves as affected by zinc deficiency. *Photosynthetica*, 43: 591-596.
- Warren J.M., Bassman J.H., Fellman J.K., Mattinson D.S. and Eigenbrode S. (2003). Ultraviolet-B radiation alters phenolic salicylate and flavonoid composition of *Populus trichocarpa* leaves. *Tree Physiology*, 23: 527-535.
- Weckx J.E.J. and Clijsters H.M.M., (1997) Zn phytotoxicity induces oxidative stress in primary leaves of *Phaseolus vulgaris*. *Plant Physiology and Biochemistry*, 35: 405-410.
- Welkie G.M., Hekmat-Shoar H. and Mille, G.W. (1990) Responses of pepper plants to iron deficiency: solution pH and riboflavin. In: Van Beusichem, M.L. (Ed.), *Plant Nutrition-Physiology and Application*. Kluwer Academic, Dordrecht, Netherlands.
- Wenrong C., Xiaoe Y., Zhenli H., Ying F. and Fenghong H. (2008) Differential changes in photosynthetic capacity, 77 K chlorophyll fluorescence and chloroplast ultrastructure between Zn-efficient and Zn-inefficient rice genotypes (*Oryza sativa*) under low zinc stress *Physiologia Plantarum*, 132: 89-101.
- Wilson K.E., Wilson M.I. and Greenberg B.M. (1998) Identification of the flavonoid glycoside that accumulates in *Brassica napus* L. cv. Topas specifically in response to ultraviolet B radiation. *Photochemistry and Photobiology*, 67: 547-553.
- Yamasaki H., Uefuji H. and Sakihama Y. (1996). Bleaching of the red anthocyanin by superoxide radical. *Archives of Biochemistry and Biophysics*, 332: 183-186.